

Comprensione dei meccanismi di azione delle sostanze biostimolanti attraverso l'integrazione della metabolomica e della trascrittomica


Prof. Luigi Lucini – Università Cattolica del Sacro Cuore
Prof. Youry Pii – Libera Università di Bolzano




2^a BIOSTIMOLANTI
CONFERENCE **DIGITAL**

Protein hydrolysate

Mixture of polypeptides, oligopeptides and small molecules that are manufactured from protein sources using partial hydrolysis



Mechanisms are still poorly understood



Rooting activity of PH




Control




PH3 foliar

Rooting activity of PH




→ Following the treatments with PH, root development showed an increasing trend


metabolomics





molecular mechanisms




Hierarchical clustering, shoots



Hierarchical clustering, roots




Supervised OPLS-DA



Rooting activity of PH

Shoot Metabolome





Metabolites involved in the synthesis of auxin are **not differentially modulated** in the leaves of treated plants as compared to untreated ones.

Rooting activity of PH

Root Metabolome

C







Rooting activity of PH

- The plant hormone auxin (IAA) is a key regulator of plant growth and development.
- Local biosynthesis and polar transport of auxin act together to generate auxin gradients.

Rooting activity of PH




Auxin concentration is tightly regulated in plants → biosynthesis vs. conjugation/degradation

Auxin-mediated signaling


Auxin Signaling

Tomasz Paciorek and Jiří Friml





© Journal of Cell Science 2006 (119), pp. 1199-1202


Aux/IAA Transcription factors



Aux/IAA Transcription factors



Auxin-mediated signaling






The Plant Cell, Vol. 17, 2676–2692, October 2005, www.plantcell.org © 2005 American Society of Plant Biologists

The Tomato *Aux/IAA* Transcription Factor *IAA9* Is Involved in Fruit Development and Leaf Morphogenesis

Hua Wang,^a Brian Jones,^a Zhengguo Li,^b Pierre Frasse,^a Corinne Delalande,^a Farid Regad,^a Salma Chaabouni,^a Alain Latché,^a Jean-Claude Pech,^a and Mondher Bouzayen^{a,1}


RT-PCR analyses









RNA-sequencing

Data analysis



Next steps



Conclusions

- ✓ Rooting experiments showed a PH-dependent stimulation of auxin response
- ✓ Distinctive effects were observed between different plant organs, despite the PHs were applied foliarly
- ✓ A broad reprogramming of metabolism was elicited by the PHs, and the metabolite signature overlapped with exogenous auxin only in shoots
- ✓ Transcription Factors involved in auxin response are activated by PH treatments; nevertheless, genome-wide transcriptomic approaches are required to gain further insight in the biological effects of PH



Fakultät für Naturwissenschaften und Technik
Facoltà di Scienze e Tecnologie
Faculty of Science and Technology



Thanks for your attention!!